





## CHEMISTRY-CY


**Q.1 – Q.30 : Carry ONE mark each.**

- The complexes  $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]\text{Br}_2$  and  $[\text{Co}(\text{NH}_3)_4\text{Br}_2]\text{Cl}\cdot\text{H}_2\text{O}$  are examples of
  - ionization isomerism
  - linkage isomerism
  - geometric isomerism
  - optical isomerism
- In the trigonal bipyramidal crystal field, the d orbital with the highest energy is
  - $d_{xy}$
  - $d_{x^2-y^2}$
  - $d_{yz}$
  - $d_z^2$
- The magnetic moment of the complex  $\text{K}_3[\text{CoF}_6]$  is  $5.0 \mu_B$ . The total stabilization energy will be
  - $-0.4 \Delta_0$
  - $-0.4 \Delta_0 + P$
  - $-2.4 \Delta_0 + 3P$
  - $-1.8 \Delta_0 + 3P$
- The metal present at the active site of the protein carboxypeptidase A is
  - Zinc
  - molybdenum
  - magnesium
  - cobalt
- The neutral complex which follows the 18-electron rule is
  - $(\eta^5 - \text{C}_5\text{H}_5)\text{Fe}(\text{CO})_2$
  - $(\eta^5 - \text{C}_5\text{H}_5)\text{Mo}(\text{CO})_3$
  - $(\eta^5 - \text{C}_5\text{H}_5)_2\text{CO}$
  - $(\eta^5 - \text{C}_5\text{H}_5)\text{Re}(\eta^6 - \text{C}_6\text{H}_6)$
- The shape of the molecule  $\text{XeO}_2\text{F}_2$  is
  - distorted tetrahedral
  - square planar
  - trigonal bipyramidal
  - tetrahedral
- Triplet superphosphate is made by treating phosphate rock with
  - conc.  $\text{H}_2\text{SO}_4$
  - conc.  $\text{HNO}_3$
  - conc.  $\text{HCl}$
  - conc.  $\text{H}_3\text{PO}_4$
- The number of hydroxy (OH) groups present in phosphorus acid is
  - one
  - two
  - three
  - four
- Out of the following, the one which is not an excitation source for IR spectrometer is
  - Tungsten filament lamp
  - Nernst glower
  - Deuterium lamp
  - Mercury arc
- One of the following is not related to polarography
  - limiting current
  - diffusion current constant
  - Ilkovic equation
  - current efficiency
- Among the following, the optically inactive compound is:







18. Order of reactivity of the following dienes X, Y and Z in Diels-Alder reaction is




- (a)  $X > Z > Y$  (b)  $Y > X > Z$   
 (c)  $Y > Z > X$  (d)  $X > Y > Z$

19. The major product P of the following reaction is



20. Among the following, the most stable isomer for 3-methoxycyclohexanol is:



21. A reaction follows second order rate law,  $-d[A]/dt = k[A]^2$ , if

- (a) a plot of  $[A]$  versus  $t$  is a straight line (b) a plot of  $1/[A]$  versus  $t$  is a straight line  
 (c) a plot of  $\ln[A]$  versus  $t$  is a straight line (d) a plot of  $e^{[A]}$  versus  $t$  is a straight line

22. In an adiabatic system, the work done to change its state from A to B is

- (a) dependent on path from A to B  
 (b) independent of path from A to B  
 (c) path dependence is related to particulars of states A and B  
 (d) path dependence is related to both states A and B and choice of path

23. The set of eigenfunctions  $\sqrt{\frac{2}{a}} \sin \frac{n\pi x}{a}$  ( $0 \leq x \leq a$ ,  $n = 1, 2, 3, \dots$ ) is

- (a) orthogonal (b) normalized  
 (c) both orthogonal and normalized (d) unnormalized

24. The function  $e^{\alpha x^2}$  ( $\alpha > 0$ ) is not an acceptable wave function for bound system, because

- (a) it is not continuous (b) it is multivalued  
 (c) it is not normalisable (d) all of these

25. First order perturbation correction  $\Delta \epsilon_n^{(1)}$  to energy level  $\epsilon_n$  of a simple harmonic oscillator due to the anharmonicity perturbation  $\gamma x^3$  is given by

- (a)  $\Delta \epsilon_n^{(1)} = \gamma$  (b)  $\Delta \epsilon_n^{(1)} = \gamma^2$  (c)  $\Delta \epsilon_n^{(1)} = \gamma^{-1}$  (d)  $\Delta \epsilon_n^{(1)} = 0$

26. Resonant frequencies for EPR and NMR are respectively in the spectral region  
 (a) microwave and far-IR (b) far-IR and microwave  
 (c) radiofrequency and microwave (d) microwave and radiofrequency
27. The  $2s$  orbital of H-atom has radial node at  $2a_0$  because  $\psi_{2s}$  is proportional to  
 (a)  $\left(\frac{1}{2} + \frac{r}{a_0}\right)$  (b)  $\left(2 + \frac{r}{a_0}\right)$  (c)  $\left(2 - \frac{r}{a_0}\right)$  (d)  $\left(2 - \frac{r}{2a_0}\right)$
28. Given that the mean speed of  $H_2$  is  $1.78 \text{ km s}^{-1}$ , the mean speed of  $D_2$  will be  
 (a)  $1.26 \text{ km s}^{-1}$  (b)  $2.52 \text{ km s}^{-1}$  (c)  $5.04 \text{ km s}^{-1}$  (d)  $3.17 \text{ km s}^{-1}$ .
29. The triple point for water is  
 (a) unique (b) depends on p but is independent of T  
 (c) depends on T but is independent of P (d) depends on both P and T
30. Hydrolysis of urea by urease is  
 (a) first order at high concentration of urea (b) zero order at high concentration of urea  
 (c) independent of the concentration of urea (d) first order with respect to both urea and urease

**Q.31 – Q.85 : Carry TWO marks each.**

31. The rate of exchange of cyanide ligands in the complexes (i)  $[\text{Ni}(\text{CN})_4]^{2-}$ , (ii)  $[\text{Mn}(\text{CN})_6]^{3-}$  and (iii)  $[\text{Cr}(\text{CN})_6]^{3-}$  by  $^{14}\text{CN}$  follow the order.  
 (a)  $\text{ii} > \text{i} > \text{iii}$  (b)  $\text{iii} > \text{i} > \text{ii}$  (c)  $\text{i} > \text{iii} > \text{ii}$  (d)  $\text{i} > \text{ii} > \text{iii}$
32. Ligand field stabilization energies are smaller for lanthanides compared to transition metals in the same oxidation state because  
 (a) size of lanthanide ions are larger (b) f orbitals interact less effectively with ligands  
 (c) size of lanthanide ions are smaller (d) lanthanides favour oxygen donor ligands
33. For the metal-olefin complexes (i)  $[\text{PtCl}_3(\text{C}_2\text{H}_4)]^-$  and (ii)  $[\text{PtCl}_3(\text{C}_2\text{F}_4)]^-$ , the correct statement is that  
 (a) carbon-carbon bond length is same both in (i) and (ii)  
 (b) carbon-carbon bond length in (i) is smaller compared to that of (ii)  
 (c) carbon-carbon bond length in (i) is larger compared to that of (ii)  
 (d) a metallacycle is formed in each complex
34. A solution containing 47 ppm of a compound X (FW 225) has a transmittance of 29.7% in a 1.5 cm cell at 400 nm. The molar absorptivity in  $\text{L mol}^{-1} \text{ cm}^{-1}$  is  
 (a)  $1.89 \times 10^3$  (b)  $1.42 \times 10^3$  (c)  $1.68 \times 10^3$  (d)  $1.79 \times 10^3$
35. The values of M–C stretching frequencies of (i)  $[\text{V}(\text{CO})_6]^-$ , (ii)  $[\text{Cr}(\text{CO})_6]$  and (iii)  $[\text{Mn}(\text{CO})_6]^+$  follow the trend  
 (a)  $(\text{ii}) > (\text{i}) > (\text{iii})$  (b)  $(\text{ii}) > (\text{iii}) > (\text{i})$  (c)  $(\text{i}) > (\text{ii}) > (\text{iii})$  (d)  $(\text{iii}) > (\text{ii}) > (\text{i})$
36. A substance undergoes a two electron reversible reduction at dropping mercury electrode, and gives a diffusion current of  $7.5 \mu\text{A}$ . When the potential at the dropping mercury electrode is  $-0.615 \text{ V}$ , the current is  $1.5 \mu\text{A}$ . The  $E_{1/2}$  (in volt) will be  
 (a)  $-0.683$  (b)  $-0.674$  (c)  $-0.652$  (d)  $-0.633$

37. The lanthanide complex (acac = acetylacetonate; phen = 1, 10-phennathroline) that do not have square antiprismatic structure is
- (a)  $[\text{Ce}(\text{NO}_3)_6]^{2-}$  (b)  $[\text{La}(\text{acac})_3(\text{H}_2\text{O})_2]$   
 (c)  $[\text{Ce}(\text{acac})_4]$  (d)  $[\text{Eu}(\text{acac})_3(\text{Phen})]$
38. Among the following, the incorrect statement about SiC is that
- (a) it is known as corundum  
 (b) it is prepared by reducing quartz with a slight excess of coke in an electric furnace at 2000–2500°C  
 (c) pure SiC is almost colourless or pale yellow  
 (d) its hardness is slightly less than diamond.
39. The incorrect statement regarding carboranes is that
- (a) carbon tends to adopt the position of the lowest coordination number on the polyhedron  
 (b) CH groups tend to be more positive than BH groups with the same coordination number  
 (c) carbon tends to keep as close as possible to other carbon atoms  
 (d) generally, arachno-carboranes are less stable thermally than the corresponding closo-carboranes
40. The incorrect statement for solid sodium chloride is that
- (a) both sodium and chloride ions adopt inert gas configuration  
 (b) the conduction band is full  
 (c) the conduction band is empty  
 (d) the valence band is full

**Q.41 – Q.48** required matching of items of *Column-I* with the appropriate items in *Column-II*. Choose the correct one from the alternatives (a), (b), (c) and (d)

41. **Column-I**

- (P) Cytochrome c  
 (Q) Calmodulin  
 (R) Chlorophyll  
 (S) Alcohol dehydrogenase

- (a) P-V, Q-VI, R-III, S-V  
 (b) P-VI, Q-III, R-IV, S-II  
 (c) R-III, R-IV, R-VI, R-III  
 (d) S-IV, S-V, S-II, S-IV

**Column-II**

- (I) Molybdenum  
 (II) Potassium  
 (III) Magnesium  
 (IV) Zinc  
 (V) Iron  
 (VI) Calcium
- (a) Q-II, Q-III, Q-IV, Q-VI  
 (b) Q-II, Q-III, Q-IV, Q-VI  
 (c) S-IV, S-V, S-II, S-IV  
 (d) S-IV, S-V, S-II, S-IV

42. **Column-1**

- P. Atomic absorption  
 Q. Chronopotentiometry  
 R. Spectrophotometry  
 S. Conductometry

**Codes:**

- (a) P-I, Q-III, R-IV, S-V  
 (b) P-VI, Q-I, R-IV, S-II  
 (c) P-II, Q-III, R-IV, S-V  
 (d) P-V, Q-VI, R-II, S-IV

**Column-2**

- I. Transition time  
 II. Cell constant  
 III. Coulomb  
 IV. Molar absorptivity  
 V. Limiting current  
 VI. Hollow cathode lamp

43. Require matching of items of **Column I** with the appropriate items in **Column - II**. Choose the correct one from the alternatives (a), (b), (c) and (d).

**Column - I**

P: Wilkinson's catalyst

Q: Speiers's catalyst

R: Water gas shift catalyst

S: Zeolite ZSM-5 catalyst

**Column - II**I. trans -IrCl(CO)(PPh<sub>3</sub>)<sub>2</sub>

II. Hydrosilylation

III. RhCl(PPh<sub>3</sub>)<sub>3</sub>

IV. Synthetic gasoline

V. hydroformylation

VI. Zinc-copper oxide.

- |       |       |      |
|-------|-------|------|
| (a)   | (b)   | (c)  |
| P-III | P-I   | P-V  |
| Q-II  | Q-V   | Q-II |
| R-VI  | R-III | R-VI |
| S-IV  | S-IV  | S-IV |

- (d)
- |       |
|-------|
| P-III |
| Q-VI  |
| R-IV  |
| S-II  |

- 44.
- Column-1**

P. Ostwald process

Q. Solvay process

R. Mond process

S. Frasch process

**Column-2**

I. Manufacture of nickel

II. Manufacture of nitric acid

III. Manufacture of Na<sub>2</sub>CO<sub>3</sub>

IV. Manufacture of silicones

V. Manufacture of caustic soda

VI. Mining of elemental sulfur

**Codes:**

(a) P-I, Q-III, R-II, S-VI

(c) P-II, Q-I, R-IV, S-V

(b) P-II, Q-III, R-I, S-VI

(d) P-III, Q-II, R-V, S-VI

- 45.
- Column-I (Compounds)**

P. cyclohexanone

Q. cyclopentanone

R. cyclobutanone

S. cyclopropanone

**Column-2 (Carbonyl stretching frequency (cm<sup>-1</sup>))**

I. 1910

II. 1715

III. 1813

IV. 1650

V. 1780

VI. 1745

**Codes:**

(a) P-I, Q-II, R-III, S-IV

(c) P-VI, Q-V, R-IV, S-III

(b) P-II, Q-VI, R-V, S-III

(d) P-I, Q-V, R-IV, S-III

- 46.
- Column-I**

P. Many electron wave function

Q. Low temperature

R. Mean speed

S. Molecular ensemble

**Column-2**

I. Adiabatic demagnetization

II. Slater determinant

III. Partition function

IV. maxwellian distribution

V. LCAO-MO

VI. Photoejection

**Codes:**

(a) P-IV, Q-I, R-VI, S-III

(c) P-II, Q-V, R-VI, S-IV

(b) P-II, Q-I, R-IV, S-III

(d) P-VI, Q-IV, R-III, S-II

47. Require matching of items of **column - I** with the appropriate items in **column - II**. Choose the correct one from the alternative (a), (b), (c) and (d).

**Column - I****Spectral Technique**

P. Rotational transition

Q. Vibrational transition

R. Electronic transition in atoms

S. Molecular ensemble

**Column - II****Selection Rule**I.  $\Delta v = \pm 1$ II.  $\Delta J = 0$ III.  $\Delta J = \pm 1$ IV.  $\Delta l = \pm 1$ V.  $\Delta m_l = \pm 1$ VI.  $\Delta v = 0$ VII.  $\Delta l = 0$ 

(a) P-I, Q-VI, R-VII, S-V

(c) P-III, Q-I, R-IV, S-V

(b) P-II, Q-I, R-IV, S-V

(d) P-I, Q-VI, R-VII, S-V

48. **Column-1 (Molecular species)**

P.  $O_2^-$ Q.  $O_2$ R.  $O_2^+$ S.  $O_2^{2-}$ **Column-2 (Bond order and magnetic property)**

I. 2.5, paramagnetic

II. 2.0, diamagnetic

III. 1.5, diamagnetic

IV. 1.0, diamagnetic

V. 2.0, paramagnetic

VI. 1.5, paramagnetic

**Codes:**


(a) P-III, Q-V, R-IV, S-III

(c) P-III, Q-V, R-IV, S-VI


(b) P-II, Q-II, R-IV, S-III

(d) P-VI, Q-V, R-I, S-IV


49. The major product P formed in the given



50. The order of reactivity towards acid catalyzed hydrolysis of the following cyclic acetals is:


(a)  $Z > Y > X$ (b)  $X > Y > Z$ (c)  $X > Z > Y$ (d)  $Z > X > Y$

51. The binaphthol (Bnp) is:




- (a) An optically active compound with (R)-configuration.  
 (b) An optically inactive compound.  
 (c) A meso compound  
 (d) An optically active compound with (S)-configuration.


52. In the given reactions, identify the correct combination of their major products P and Q [LDA =  $\text{LiN}(\text{i-Pr})_2$ ]



53. The major stereoisomer obtained in the reaction of (S)-2-phenylpropanal with  $\text{MeMgBr}$  is:





54. The major product P formed in the following reaction is








55. Iodo-lactonization of  $\beta, \gamma$ -unsaturated carboxylic acid X with  $I_2$  and  $NaHCO_3$  gives.




56. The major stereoisomer P obtained in the following reaction is:



57. The major product P of the following reaction is:





58. Cis- and trans-2-methyl-5-*t*-butyl-1,3-dioxane each can exist as two conformers as shown below





The preferred conformations for the cis- and trans-compounds will be

- (a) P, R (b) Q, S (c) P, S (d) Q, R

59. The major product P of the given reaction is



60. The major product P formed in the following photochemical reaction is




61. An organic compound having molecular formula  $\text{C}_8\text{H}_{12}\text{O}_2$  exhibits the following peaks in IR and  $^1\text{H}$  NMR spectra.

IR :  $1720 \text{ (cm}^{-1}\text{)}$

$^1\text{H}$  NMR :  $6.95 \text{ (1H, d, } J = 8.5 \text{ Hz)}$ ,  $5.90 \text{ (1H, d, } J = 8.5 \text{ Hz)}$ ,  $4.53 \text{ (1H, q, } J = 6 \text{ Hz)}$ ,

$1.41 \text{ (3H, d, } J = 6 \text{ Hz)}$ ,  $1.20 \text{ (3H, s)}$ ,  $1.15 \text{ (3H, s)}$



62. The phase diagram of  $\text{NaCl}-\text{H}_2\text{O}$  is of simple eutectic type. The eutectic composition is 23.3 weight%  $\text{NaCl}$  and it freezes at  $-21.1^\circ\text{C}$ . The phases present in a solution containing 10 weight%  $\text{NaCl}$  at  $-20^\circ\text{C}$  are  
 (a) ice +  $\text{NaCl}$  solution (b) ice + solid  $\text{NaCl}$   
 (c)  $\text{NaCl}$  + pure water (d)  $\text{NaCl}$  + saturated  $\text{NaCl}$  solution
63. Hydrogen adsorption on a platinum surface is  
 (a) Endothermic with positive  $\Delta S$  and positive  $\Delta G$   
 (b) Endothermic with positive  $\Delta S$  and negative  $\Delta G$   
 (c) Exothermic with negative  $\Delta S$  and negative  $\Delta G$   
 (d) Exothermic with positive  $\Delta S$  and negative  $\Delta G$
64. In the reversible chemical reaction taking place under standard condition at 298 K and 1 atm in a Daniel cell,  
 $\text{Zn} | \text{Zn}^{2+}(\text{aq}) || \text{Cu}^{2+}(\text{aq}) | \text{Cu}$   
 the heat change is:  
 (a) equal to  $\Delta H^0$  (b) equal to  $T\Delta S^0$  (c) equal to zero (d) equal to  $\Delta U^0$
65. The orbital  $\psi = 1s_{\text{H}_A} - 1s_{\text{H}_B}$  of water belongs to the irreducible representation  
 (a)  $A_1$  (b)  $B_1$  (c)  $A_2$  (d)  $B_2$
66. The vibrational partition function for a molecule with fundamental frequency  $\nu$  is given by  
 (a)  $\exp\left(-\frac{\hbar\nu}{k_B T}\right)$  (b)  $\left[1 - \exp\left(-\frac{\hbar\nu}{k_B T}\right)\right]^{-1}$   
 (c)  $\exp\left(-\frac{\hbar\nu}{k_B T}\right) \left[1 - \exp\left(-\frac{\hbar\nu}{k_B T}\right)\right]^{-1}$  (d)  $\exp\left(-\frac{\hbar\nu}{2k_B T}\right) \left[1 - \exp\left(-\frac{\hbar\nu}{k_B T}\right)\right]^{-1}$
67. The internal pressure,  $\pi_T = T\left(\frac{\partial P}{\partial T}\right)_V - P$  for one mole a Vander waals gas is  
 (a)  $\frac{a}{V^2}$  (b)  $\frac{a}{V^2} \left(\frac{RT}{V-b}\right)$  (c) Zero (d)  $\frac{RT}{V-b}$
68. A gaseous sample on exposure to total radiant energy of 6.626 J at 300 nm results in the photodissociation of  $10^{-3}$  mol of this sample. Assuming the sample absorbs all the light, the quantum yield for this photochemical reaction is:  
 (a) 6.023 (b) 0.602 (c) 60.230 (d) 0.060
69. If standard emf of the cell,  
 $\text{Cu} | \text{Cu}^{2+}(\text{aq}) || [\text{Cu}(\text{NH}_3)_4]^{2+}, \text{aq. NH}_3 | \text{Cu}$   
 is 0.35, then stability constant of the formation cupric amine complex is  
 (a)  $1.0 \times 10^{27}$  (b)  $8.4 \times 10^5$  (c)  $7.0 \times 10^{11}$  (d)  $4.3 \times 10^{13}$
70. Standard entropy of crystalline carbon monoxide (in J/mol) at 0 K is around  
 (a) 0.03 (b) 2.50 (c) Zero (d) 5.76
71. Metals used in automobile catalytic converters are:  
 (a) Pt and Pd (b) Pt and Rh (c) Pd and Rh (d) Rh and Ni

**Q. 72 to Q. 77** contain a Statement with a Reason and an Assertion. For each question, choose the correct answer from the following four choices.

- (a) Both Reason and Assertion are correct      (b) Both Reason and Assertion are wrong  
(c) Reason is correct but Assertion is wrong      (d) Reason is wrong but Assertion is correct

72. **Statement:** The characteristic spectroscopic feature of the quadruply bonded  $[\text{Re}_2\text{Cl}_8]^{2-}$  is a strong royal blue colour

**Reason:** This is due to an absorption band in the visible region due to excitation of an electron from  $\sigma^2\pi^4\delta^2$  ground state to  $\sigma^2\pi^4\delta^1\delta^{*1}$  excited state

**Assertion:** This transition is quantum mechanically allowed

73. **Statement:** For the reaction  $\text{L}_n\text{MH} \rightarrow \text{L}_n\text{M}^- + \text{H}^+$ , the important factors are the strength of the M–H bond and the nature of the ligand, L

**Reason:** The key here is the stability of the complex ion,  $\text{L}_n\text{M}^-$

**Assertion:** Weak  $\pi$ -bonding ligands will stabilize  $\text{L}_n\text{M}^-$  and so will disfavour the forward reaction.

74. **Statement:** D-Glucose and D-mannose give the same phenylosazone.      [GATE 2005]

**Reason:** Osazone formation results in a loss of the stereocentre at  $\text{C}_2$  but does not affect other stereocenters.

**Assertion:** D-Glucose and D-mannose are enantiomers.

- (a) Both **Reason** and **Assertion** are correct  
(b) Both **Reason** and **Assertion** are wrong.  
(c) **Reason** is correct but **Assertion** is wrong.  
(d) **Reason** is wrong but **Assertion** is correct.

75. **Statement:** Nucleosides are stable in dilute base but undergo hydrolysis in dilute acid.      [GATE 2005]

**Reason:** Nucleosides have an N-glycosidic linkage.

**Assertion:** N-Glycosidic linkage behaves like an O-glycosidic linkage which is rapidly hydrolyzed by aqueous acid but stable in aqueous base.

- (a) Both **Reason** and **Assertion** are correct  
(b) Both **Reason** and **Assertion** are wrong.  
(c) **Reason** is correct but **Assertion** is wrong.  
(d) **Reason** is wrong but **Assertion** is correct.

76. **Statement :** For the reaction of  $\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$  the rate constant is observed to decrease with temperature.

**Reason :** As per the proposed mechanism, the first step is the dimerization of nitric oxide which is exothermic.

**Assertion :** Rate law =  $k_2K_1[\text{NO}]_2[\text{O}_2]$

- (a) Both Reason and Assertion are correct  
(b) Both Reason and Assertion are wrong  
(c) Reason is correct but Assertion is wrong  
(d) Reason is wrong but Assertion is correct.

77. **Statement:** Hydrogen gas gets warmer on expanding under isenthalpic condition

**Reason:** Joule Thomson coefficient for hydrogen is  $-0.03 \text{ K/atm}$

**Assertion:** Attractive forces are the dominant intermolecular interactions in hydrogen gas at 273 K.

**Common Data for Q. 78, Q.79 and Q.80:**

Vapour pressures of water above pure liquid water 24, 529 and 760 torr respectively at 298, 363 and 373 K. Use these data to answer the questions 78, 79 and 80.

78. Change in chemical potential (in kJ/mol) for the equilibrium  $\text{H}_2\text{O}(\text{liquid}) = \text{H}_2\text{O}(\text{gas})$  at 298K is

- (a) 8.6      (b)  $-3.8$       (c) 7.87      (d) 3.72

79. Aqueous solution of sodium chloride ( $\chi_{\text{NaCl}} = 0.015$ ) at 298 K is in equilibrium with a water vapour pressure (in torr) of  
 (a) 23.64 (b) 748.60 (c) 24.36 (d) negligible
80. Average value of enthalpy of vaporisation (in kJ/mol) of water between 363 and 373 K is  
 (a) 42.50 (b) 40.80 (c) -40.65 (d) -40.80

**Linked Answer Q.81(a) and Q.81(b):**

- 81.(a) As per Huckel theory,  $\pi$ -electron energy levels of cyclobutadiene are  
 (a)  $\alpha + 2\beta, \alpha + \beta, \alpha - \beta, \alpha - 2\beta$  (b)  $\alpha + 2\beta, \alpha - \beta, \alpha - \beta, \alpha - 2\beta$   
 (c)  $\alpha + 2\beta, \alpha, \alpha, \alpha - 2\beta$  (d)  $\alpha + \beta, \alpha - \beta, \alpha - \beta, \alpha - 2\beta$
- 81.(b) Given that  $\beta = -75 \text{ kJ/mol}$ , cyclobutadiene is  
 (a) paramagnetic and its lowest absorption energy is 150 kJ  
 (b) paramagnetic and its lowest absorption energy is 75 kJ  
 (c) diamagnetic and its lowest absorption energy is 75 kJ  
 (d) diamagnetic and its lowest absorption energy is 150 kJ.

**Linked Answer Q.82(a) and Q.82(b):**


- 82.(a) For the complex ion  $[\text{Cu}(\text{NH}_3)_6]^{2+}$ , the coordination geometric will be  
 (a) octahedral (b) tetragonally distorted octahedral  
 (c) trigonal prismatic (d) trigonal antiprismatic
- 82.(b) The number of possible d-d transitions will be  
 (a) one (b) two (c) three (d) four



**Linked Answer Q.83(a) and Q.83(b):**

- 83.(a). The following data was obtained with the GLC. Column temperature,  $60^\circ\text{C}$ , inlet pressure, 1270 torr, outlet pressure, 770 torr, flow rate of carrier gas at  $25^\circ\text{C}$ , 18 mL/min and retention time for air, 0.30 min, the pressure drop correction factor will be  
 (a) 0.648 (b) 0.740 (c) 0.770 (d) 0.715
- 83.(b). Corrected retention volume for air (mL) will be  
 (a) 4.02 (b) 4.72 (c) 4.46 (d) 4.25





**Linked Answer Q.84(a) and Q.84(b):**

- 84.(a). The major product P of the following reaction is:






84.(b). Major compound Q obtained on reaction of P with NaH in DMF is:




**Linked Answer Q.85(a) and Q.85(b):**

85. (a) In the following sequence of reactions, the major product Q is:



85.(b) The major product on sulphonation of Q with  $\text{H}_2\text{SO}_4$  at  $160^\circ\text{C}$  is:



\*\*\*\*\* END OF THE QUESTION PAPER \*\*\*\*\*