PAPER : CSIR-UGC-NET/JRF DEC. 2024

CHEMICAL SCIENCES

PART-B

- 21. Nano Particles have length scales
 - (a) between 0.1 to 1000 nm in any dimension
 - (b) in any nanometer scale and always in one particular dimension
 - (c) between 1 to 500 nm and always in one particular dimension
 - (d) between 1 to 100 nm in any dimension
- 22. The correct order for the X-H bond dissociation energies (BDE) in the following compounds is
 - (a) $Me_3Si H > Me_3C H > Me_3Sn H$
 - (b) $Me_3C-H > Me_3Si-H > Me_3Sn-H$
 - (c) $Me_3C-H > Me_3Sn-H > Me_3Si-H$
 - (d) $Me_3Sn H > Me_3C H > Me_3Si H$
- 23. The correct option for the pKa of following pairs of compounds is



 $B_2H_6 + NH_2CH_3 \longrightarrow A$

24.

```
B_2H_6 + NMe_3 \longrightarrow B
```

- (a) H₃BNH₂CH₃ and H₃BNMe₃
- (b) $[BH_2(NH_2CH_3)_2]^+[BH_4]^-$ and H_3BNMe_3
- (c) $H_3BNH_2CH_3$ and $[BH_2(NMe_3)_2]^+[BH_4]^-$
- (d) $[BH_2(NH_2CH_3)_2]^+[BH_4]^-$ and $[BH_2(NMe_3)_2]^+[BH_4]^-$

25. The co-enzymes involved in the following biosynthesis are





26. For the reaction of RBr with EtONa/EtOH, the correct match for the R groups in **Column P** with the relative rates in **Column Q** is (for $R = CH_3CH_2$, relative rate = 1)

	Column - P		Column - Q
A.	CH ₃	i.	4.2×10^{-6}
B.	CH ₃ CH ₂ CH ₂	ii.	0.03
C.	$(CH_3)_2 CHCH_2$	iii.	0.28
D.	$(CH_3)_3 CCH_2$	iv.	17

- (a) A-iv; B-iii; C-ii; D-i
- (c) A-iii; B-ii; C-iv; D-i

(b) A-iv; B-i; C-iii; D-ii(d) A-ii; B-iii: C-i; D-iv

- 27. The option showing the correct structural types for ZnFe₂O₄ and KMnF₃, respectively, is
 (a) Perovskite and Fluorite
 (b) Perovskite and Antifluorite
 (c) Spinel and Perovskite
 (d) Spinel and Fluorite
- 28. In a 3-dimensional isotropic harmonic oscillator, the degeneracy of the state with energy equal to (9/2) $\hbar \omega$ is [ω is the angular frequency] (a) 3 (b) 9 (c) 6 (d) 10
- 29. The following reaction occurs under



- (a) photochemical conditions via a disrotatory ring-opening
- (b) photochemical conditions via a conrotatory ring-opening
- (c) thermal conditions via a disrotatory ring-opening
- (d) thermal conditions via a conrotatory ring-opening

30. The order of acceptor strength towards Me₂S donor is

(a)
$$AlCl_3 > BCI_3 > GaCl_3$$
(b) $BCI_3 > GaCl_3 > AICI_3$ (c) $GaCl_3 > AICI_3 > BCI_3$ (d) $AlCl_3 > GaCl_3 > BCI_3$

31. The number of D and P terms that arise from p³ electronic configuration of an atom, respectively, are

- (a) 3 and 2 (b) 2 and 3 (c) 2 and 2 (d) 1 and 1
- 32. The solvent that shifts the Schlenk equilibrium to the right side is

 $2RMgX \Longrightarrow MgR_2 + MgX_2$

(R = isopropyl)(a) hexane(b) tetrahydrofuran(c) dioxane(d) diethyl ether



- 33. The standard reduction potentials of lanthanides (Ln³⁺/Ln) are
 - (a) similar to each other and also similar to those of late transition metals
 - (b) different from each other but similar to lighter p-block elements
 - (c) similar to each other and also similar to those of s-block elements
 - (d) different from each other but similar to those of s-block elements
- 34. The correct order for the relative rate of addition of *CCl₃ to 2-methylpropene (A), styrene (B) and 2-methylbut-2-ene (C) is
 - (a) A > B > C(b) B > A > C(c) $\mathbf{B} > \mathbf{C} > \mathbf{A}$ (d) $\mathbf{C} > \mathbf{B} > \mathbf{A}$
- The ionic strength (in mol kg⁻¹) of an aqueous solution of 0.03 mol kg⁻¹ K₃[Fe(CN)₆] is closest 35. to
 - (a) 0.27 (b) 0.18 (c) 0.12(d) 0.15
- Consider the statements for the complexes $[RhCl_3(H_2O)_3]$ (X) and $[Ir(CO)(Cl)(PPH_3)_2]$ (Y) 36. (A) X has two isomers
 - (B) Y has two isomers
 - (C) Both X and Y are prone to oxidative addition
 - (D) d_{2} orbital is most destabilized in Y

The correct option is

- (a) A, B and C only (b) B, C and D only (c) A and B only (d) A and C only
- 37. Quantum confinement leads to
 - (a) increase in the band gap of the semiconductors
 - (b) decrease in the band gap of the metal nanoparticles
 - (c) decrease in the band gap of the semiconductors
 - (d) no change in the band gap of the quantum dots
- 38. For a given metal ion, the correct order of the nephelauxetic effect of ligands is (en = ethylene diamine) CAREER END (b) $I^- > CN^- > NH_3 > en$
 - (a) $CN^{-} > en > NH_{3} > I^{-}$
 - (c) $CN^{-} > I^{-} > en > NH_{3}$

- (d) $I^- > CN^- > en > NH_3$
- The enzyme nitrogenase converts one mole of N_2 to x mole of NH_3 and y mole of H, using z mole 39. of protons and w mole of electrons. The values of x, y, z, and w, respectively, are (c) 1, 2, 8, and 8 (d) 2, 1, 8, and 8 (a) 2, 0, 6, and 6 (b) 1, 0, 6, and 6
- 40. Correct statement about polymerization is
 - (a) The average molar mass of the polymer product does not depend upon the time length of stepwise polymerization
 - (b) The slower the initiation of the chain, the higher the average molar mass of the polymer in chain polymerization
 - (c) In chain polymerization, an activated monomer attacks a minimum of three other monomers to link
 - (d) The average chain length of a polymer in stepwise polymerization is linearly dependent on the fraction of the reacted monomers



41. The steps involved in the reaction of acetaldehyde with formaldehyde in the presence of NaOH to produce pentaerythritol $[C(CH_2OH)_4]$ are (a) Claisen condensation followed by Knoevenagel condensation (b) Cannizzaro reaction followed by Claisen condensation (c) Knoevenagel condensation followed by aldol reactions (d) aldol reactions followed by Cannizzaro reaction 42. The lowest energy π -MO of butadiene has an energy of [β is resonance energy] (c) 0.61804 β (a) -1.61804β (b) -0.61804β (d) 1.61804 β The radii of the cation and anion of an ionic compound are 74 pm and 170 pm, respectively. The 43. coordination number of the cation and the best possible geometry of the compound are, respectively (c) 8, cubic (a) 4, tetrahedral (b) 6, octahedral (d) 4, square planar 44. The half-lives for the forward and reverse reactions that are first order in both directions, are 24 ms and 39 ms, respectively. The relaxation time for return to equilibrium after a temperature jump is closest to (a) 21 ms (b) 32 ms (c) 43 ms (d) 11 ms 45. At T = 0K, the entropy (in JK⁻¹) of 2 moles of CO is closest to (b) 5.76 (a) 0(c) 11.53 (d) 23.05 46. For an electron in a hydrogen atom, with azimuthal quantum number, l = 1 and magnetic quantum number, m = 1, the angle (in degrees) between the z-axis and the orbital angular momentum vctor is (a) 0(b) 45 (c) 54.7 (d) 90 47. The correct statement is (en = ethylene diamine)(a) K₂[CuF_c] is paramagnetic and sodium nitroprusside is diamagnetic (b) Both $K[AgF_4]$ and $K_3[CuF_6]$ are diamagnetic (c) Both $[Co(en)_3]Cl_3$ and $K_3[CuF_6]$ are paramagnetic (d) Sodium nitroprusside is paramagnetic and [Co(en),]Cl, is diamagnetic 48. The major product formed in the following reaction is H_2 (1 atm) Lindlar's catalyst SiMe₃ Me quinoline, hexane o °C SiMea (b) Me SiMe₃ (d) Me SiMes



49. In the ¹H NMR, the methylene protons of ClCH₂-C(Cl)(Br)CH₃ (**X**) and CH₃CH₂-C(Cl₂)CH₃ (**Y**) appear as (a) X=Y=AB quartet (b) **X**=**Y**=quartet (c) **X**=AB quartet; **Y**=quartet (d) X=quartet; Y=AB quartet 50. Consider the statements (A) Decomposition of H_2O_2 in aqueous solution catalyzed by bromide ion is a homogeneous catalytic reaction. (B) Hydrogeneration of ethene to ethane accelerated by Pd or Ni particles is a heterogeneous catalytic reaction. (C) Enzymes increase the equilibrium constants of the reactions. (D) Turnover number is the number of catalytic cycles till the catalyst becomes inactive The set of correct statements is (a) A, B and C only (b) B, C and D only (c) A, B and D only (d) C, D and A only A two-level system consists of a double degenerate excited state which is \in energy above the 51. ground state. The y-intercept of the $ln(P_e/P_g)$ vs 1/T plot is (P_e and P_g are the probabilities associated with the excited and ground states, respsectively.) (a) 0 (b) 2(c) 2 ln 2 (d) ln 2 In the absence of nitrogen, the enzyme nitrogenase functions as 52. (a) nitrile hydratase (b) carboxypeptidase (c) urease (d) hydrogenase 53. The following structures are Et and Ėt Ξ (b) enantiomers AREER END (a) identical (d) constitutional isomers (c) diastereomers 54. The rotational quantum number associated with the most intense transition in the microwave spectrum of a diatomic molecule varies with temperature (T) as (b) \sqrt{T} (c) T^2 (a) T (d) $1/\sqrt{T}$ The order of polarity of the B-C bond for the following compounds 55. $B(C = CH)_{3}(P); B(CH = CH_{2})_{3}(Q); B(CH_{2} - CH_{3})_{3}(R); B(CH_{2} - C_{6}H_{5})_{3}(S)$

(a)
$$R < S < Q < P$$
 (b) $S < R < Q < P$ (c) $R < S < P < Q$ (d) $P < Q < R < S$



56. The reaction of dipeptide, H_2N -Gly-Ala-CO₂H, with PhNCS at pH 8 followed by treatment with CF₃COOH produces



60. For the following equilibrium, the correct match for the carbonyl compounds in **Column P** with the equilibrium constant K in **Column Q** is



	Column-P		Column - Q
А.	CH ₃ CHO	i.	1.2×10^{6}
В.	Cl ₃ CCHO	ii.	1.06
C.	$CF_3C(O)CF_3$	iii.	2000

(a) A-iii; B-ii; C-i

(b) A-ii; B-iii; C-i

(c) A-ii; B-i; C-iii (d) A-i; B-iii; C-ii



PART-C

61. A correct statement, which <u>always</u> holds good, involving the zeroth-order (E_0^0) , first-order (E_0^1)

and second-order (E_0^2) perturbed energies for the ground state is $[E_0]$ is the exact ground state energy]

- (a) $E_0^0 > E_0^1 > E_0^2 > E_0$ (b) $E_0^0 > E_0^1 > 0$ (c) $E_0^0 + E_0^1 \ge E_0$ (d) $E_0^2 > 0$
- 62. The conformation of the reactant that gives (E)-1, 2-diphenyl-1-propene in the following reaction is



- 63. The expression for d-orbitals with n = 3, l = 2 and m = ±2 is $\psi_{32\pm 2} = NR'(r)r^2 \sin^2 \theta e^{\pm 2i\phi}$ where N is a constant, r, θ , ϕ are spherical polar co-ordinate. R'(r) is a function of r. The orbital generated from a linear combination of ψ_{322} and ψ_{32-2} orbitals, $\frac{1}{i}(\psi_{322} \psi_{32-2})$, is
 - (a) d_{z^2} (b) d_{xy} (c) d_{yz} (d) d_{zx}
- 64. The correct set of reagents to effect the following transformation is



- (a) TEMPO, NCS, CH_2Cl_2/H_2O (pH 8.6)
- (b) MnO_2 , acetone
- (c) CrO_3 , H_2SO_4 , H_2O -acetone
- (d) TEMPO (cat.), NaOCl (cat.), NaClO₂, toluene/phosphate buffer pH 6.8



65. The major product formed in the following reaction is





66. The ground state term, Lande factor (g), and the calculated magnetic moment (m_{calc}) for Pr^{3+} (atomic number = 59) are (a) ${}^{3}H_{4}$, 0.80, 2.68 (b) ${}^{3}H_{4}$, 0.80, 3.58 (c) ${}^{5}I_{4}$, 1.33, 2.68 (d) ${}^{6}H5/2$, 1.33, 3.58

67. Match the correct set of IR bands to the given compounds ($Cp = C_5H_5$, $Cp^* = C_5Me_5$)

$(a) Cp_2 Ti (CO)_2$	(i) 1979 and 1897 cm^{-1}
$(b) CpCp^*Ti(CO)_2$	(ii) 1956 and 1875 cm^{-1}
$(c) Cp_2^* Ti (CO)_2$	(iii) 1930 and 1850 cm^{-1}

(a) a-ii, b-iii, c-i

(b) a-iii, b-i, c-ii (c) a-i, b-iii, c-ii

, c-ii (d) a-i, b-ii, c-iii

68. The major product formed in the following reaction is



69. One mole of a monoatomic ideal gas at 1 atm pressure undergoes compression from 49.2 L to 24.6 L under adiabatic reversible conditions. The final temperature (in K) of the gas is closest to (a) 952 (b) 848 (c) 756 (d) 1697



73.

70. For the following transformation, the product is formed through



- (a) an S_N^2 reaction
- (b) an S_{N1} reaction
- (c) a 1,2-elimination followed by a 1,4-addition reaction
- (d) a 1,4-elimination followed by a 1,4-addition reaction
- 71. A chromium carbonyl compound $Cr(CO)_6$ reacts with NaBH₄ to give **A**. The Lewis base **A** reacts with another molecule of $Cr(CO)_6$ to form compound **B** with the release of CO. In another reaction, compound **A** reacts with BH₃ to produce **C**. Compounds **A**, **B** and **C**, respectively, are (a) $[Cr(CO)_5(BH_4)]^-$, $[(CO)_5Cr(BH_4)-Cr(CO)_5]^-$ and $[Cr(CO)_4B_2H_7]^-$
 - (b) $[Cr(CO)_5H]^-$, $[(CO)_5Cr-H-Cr(CO)_5]^-$ and $[Cr(CO)_4BH_4]^-$
 - (c) $[Cr(CO)_{5}(BH_{4})]^{-}$, $[(CO)_{5}Cr-BH_{4}-Cr(CO)_{5}]^{-}$ and $[Cr(CO)_{5}BH_{4}]^{-}$
 - (d) $[Cr(CO)_5H]^-$, $[(CO)_4Cr-H-Cr(CO)_6]^-$ and $[Cr(CO)_5BH_4]^-$
- 72. The correct option of the isomer Shifts in ¹¹⁹Sn Mossbauer spectra for the following compounds is



- 74. Consider the following reactions,
 - A. ${}^{210}_{84}\text{Po} \rightarrow {}^{206}_{82}\text{Pb} + P$ B. $\left[{}^{125}_{52}\text{Te}\right]^* \rightarrow {}^{125}_{52}\text{Te} + Q$ C. ${}^{14}_{6}\text{C} \rightarrow {}^{14}_{7}\text{N} + R$ D. ${}^{23}_{12}\text{Mg} \rightarrow {}^{23}_{11}\text{Na} + S$ The correct option for **P**, **Q**, **R** and **S** is (a) $P = {}^{4}_{2}\text{He}$, $Q = \gamma$, $R = {}^{0}_{-1}\text{e}$, $S = {}^{0}_{1}\text{e}$



(c) $P = {}^{0}_{-1}e$, $Q = {}^{0}_{1}e$, $R = {}^{4}_{2}He$, $S = \gamma$

(d) $P = {}_{2}^{4}He$, $Q = \gamma$, $R = {}_{1}^{0}e$, $S = {}_{-1}^{0}e$

75. The correct statement about the following reaction is



- (a) A is formed as major product via a chair-like transition state
- (b) B is formed as major product via a chair-like transition state
- (c) A is formed as major product via a boat-like transition state
- (d) B is formed as major product via a boat-like transition state
- 76. A particle, confined in a one-dimensional box between x = 0 to x = L is perturbed by a constant potential V on the left half of the box (x = 0 to x = L / 2) and by V/3 on the right half (x = L / 2 to x = L). The first-order perturbation correction to the ground state energy is (a) V/2 (b) 2V/3 (c) 3V/4 (d) 3V/2
- 77. The average time for which a hydrogen atom remains adsorbed on a given surface is 35% shorter at 1000 K than at 600 K. The activation energy (in kJ mol⁻¹) for desorption is closest to
 (a) 2.3
 (b) 3.4
 (c) 4.5
 (d) 5.4

78. Mark-Houwink equation can be used to determine molecular weight of a polymer. The values of empirical constants are 1.6×10^{-4} dL g⁻¹ and 0.60. If the intrinsic viscosity of the polymer solution is 0.04 dL g⁻¹, the molar mass (in g mol⁻¹) of the polymer is closest to (a) 10000 (b) 1101 control (c) 16000 (d) 9600

- 79.If mean of a data set (25, 29, 25, 32, 24 and x) is 27, then the median is(a) 32(b) 27(c) 26(d) 25
- 80. The correct catalyst for conversion of acetophenone to (S)-1-phenylethanol under high pressure of hydrogen is
 - [DPEN = 1,2-diphenyl-1,2-ethylenediamine]
 - (a) [(S)-BINAP]RuCl₂
 - (b) [(R)-BINAP]RuCl₂
 - (c) [(S)-BINAP][(S,S)-DPEN]RuCl₂/t-BuOK
 - (d) [(R)-BINAP][(R,R)-DPEN]RuCl₇/t-BuOK

81. In a diffusion-controlled reaction in benzene between two species with similar radii having 2.0 mol m⁻³ initial concentrations, the time (in ns) required for the concentration of the species to fall to half of their initial values at 320 K is closest to

[The viscosity coefficient of benzene is 0.601 cP and 1 cP= 10^{-3} kg m⁻¹s⁻¹] (a) 26 (b) 76 (c) 42 (d) 62



82. Complex







The option showing complexes with their correct electronic configuration and bond order is

- (a) a-iii-q; b-i-p; c-ii-r
- (b) a-i-p; b-iii-r; c-ii-q(d) a-ii-q; b-i-r; c-iii-p
- (c) a-ii-r; b-iii-q; c-i-p
- 83. In the extraction of lanthanides, when an aqueous solution of Ln^{3+} is poured into a cation exchange resin column, the Ln^{3+} that moves fastest through the resin is
 - (a) Lu^{3+} (b) La^{3+} (c) Gd^{3+} (d) Sm^{3+}



84. The intermediate involved in the following transformation is



85. The correct reagent to effect the following transformation is



- (a) 6N HCl
 (b) 1N NaOH
 (c) Morpholine, Pd(PPh₃)₄
 (d) CF₃CO₂H
 86. At 0 °C, the standard volume of transition from ice to water is -1.6 cm³ mol⁻¹ and the corresponding standard entropy of transition is 22 JK⁻¹mol⁻¹. An increase in pressure by 100 bar would result in lowering of freezing point (in K) of water by (1 bar=10⁵ Pa)
 (a) 1.23
 (b) 0.56
 (c) 0.73
 (d) 1.46
- 87. In the presence of single electron transfer reagent, SmI_2 , the major product formed in the following reaction is



- 88. The total number electronic transitions between triplet D and triplet F multiplets due to spin-orbit coupling is
 - (a) 6 (b) 5 (c) 3 (d) 1
- 89. Consider the reactions in List I and related enzymes in List II

List I	List II
superoxide to oxygen	amine oxidase
hydrolysis of peptide	Ni-superoxide dismutase
hydroxylation of camphor	carboxypeptidase
primary amine to aldehyde	cytochrome P450
The option showing the correct match is	
(a) a-ii, b-iii, c-iv, d-i	(b) a-iv, b-i, c-ii, d-iii
(c) a-iii, b-ii, c-i, d-iv	(d) a-ii, b-i, c-iv, d-iii



90. The major product formed in the following reaction is $\frac{OH}{MeNO_2, 25^{\circ}C}$









C _{2v}	E	C ₂	σ_{v}	σ'_v			
A ₁	1	1	1	1	z	x^2, y^2, z^2	
A ₂	1	1	-1	-1	R _z	xy	
B ₁	1	-1	1	-1	x,R _y	yz	
B ₂	1	-1	-1	1	y, R _x	ZX	

91.

For formaldehyde (character table is shown above), the allowed electronic transition by x-polarized light is

(a)
$${}^{1}A_{1} \rightarrow {}^{1}A_{1}$$
 (b) ${}^{1}A_{1} \rightarrow {}^{1}A_{2}$ (c) ${}^{1}A_{1} \rightarrow {}^{1}B_{1}$ (d) ${}^{1}A_{1} \rightarrow {}^{1}B_{2}$

92. The major product formed in the following reaction is



- 93. Colloidal solutions are stabilized by
 - (a) van der Waal's forces
 - (b) small particle size
 - (c) shape of particles
 - (d) electrical double layer at the surface of the particles



94. The major product formed in the following reaction is



95. The most stable conformation of the following compound is



- 96. Consider the following statements regarding the magnetic properties of lanthanide ions.
 - A. Observed magnetic moments are highly dependent on the ligand field

(b) B and C only

B. Only ground J state is populated

C. Spin-orbit couplings are in the order of ~1000 cm⁻¹ while the ligand field effects are only about ~100 cm⁻¹

D. The spin-only formula cannot be used to calculate the magnetic moment of f^7 configuration The option showing the correct statements is

(a) A and B only

- (c) A and D only (d) C and D only
- 97. The major product formed in the following reaction is







98. Given are the statements regarding the overall stability constants $(\log \beta)$ for the formation of $[M(en)_3]^{2+}$ and $[M (EDTA)]^{2-}$ (en = ethylene diamine, EDTA = ethylenediamine tetraacetate), where M^{2+} is a divalent metal ion ($M^{2+} = Mn^{2+}$) F^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} , Zn^{2+})

A. The $(\log \beta)$ is lowest for Mn²⁺ in both $[M(en)_3]^{2+}$ and $[M(EDTA)]^{2-}$ series $(Mn^{2+} \text{ to } Zn^{2+})^{2-}$

B. The $(\log \beta)$ value for $[Mn(EDTA)]^{2-}$ is lower than $[Mn(en)_{3}]^{2+}$

C. The $(\log \beta)$ values increase in the series $(Mn^{2+} \text{ to } Zn^{2+})$ for both EDTA and "en" complexes

(b) A, B and D only

D. ΔS^0 remains nearly constant along the series

- The option with the correct statements is
- (a) B and C only
- (c) A and D only (d) A, B and C only

99. The atomic mass of X and Y are 5 amu and 40 amu, respectively. For the diatomic molecule XY, the spacing between any two successive lines is 8 cm^{-1} in the microwave spectrum. The bond

length of XY (in overline A) is closest to
$$\left(\frac{h}{8\pi^2 c} = 2.8 \times 10^{-44} \text{ Js}^2 \text{m}^{-1}, 1 \text{ amu} = 1.667 \times 10^{-27} \text{ kg}\right)$$

(a) 0.688 (b) 0.974 (c) 1.377 (d) 1.948

100. The fundamental vibrational frequencies of H₂ and H³⁷Cl are 4395 cm⁻¹ and 2988 cm⁻¹ respectively. Considering all molecules are in their respective ground vibrational state, the energy change (in cm⁻¹) of the reaction below

$$HD + H^{37}Cl \rightarrow H_2 + D^{37}Cl$$

is closest to [Assume that force constant remains same with isotopic substitution] (a) -65 (b) -130 (c) -260 (d) -520

101. For the given cell,

$$Zn(s)|Zn^{2+}(aq, 0.5 M)||Ag^{+}(aq, 0.1 M)|Ag(s)|$$

the emf (in V) of the cell at 25 °C is closest to

[At 25°C,
$$E_{Zn^{2+}/Zn}^{0} = -0.76 \text{ V}$$
 and $E_{Ag^{+}/Ag}^{0} = +0.80 \text{ V}$
(a) 0.05 (b) 0.04 (c) 1.56 (d) 1.51





$\begin{array}{c} \text{Enzymes} \rightarrow \\ \text{Moieties} \downarrow \end{array}$	Haemoglobin	Nickel-superoxide dismutase	[FeFe] hydrogenase	Hemerythrin
А	tyrosine	µ-hydroxo	histidine	aza (dithiolato)
В	histidine	µ-hydroxo	aza (dithiolato)	tyrosine
С	histidine	A P = tyrosine	aza (dithiolato)	µ-hydroxo
D	µ-hydroxo	aza(dithiolato)	histidine	tyrosine

The correct option is

(a) A

A (b) B

(c) C

(d) D

105. The major product formed in the following reaction sequence is







109.

106. A particle incident from the region x < 0 is under a potential barrier with finite height V₀, and finite width *L*, as in the diagrams. When the total energy (*E*) of the incident particle is less than $\begin{pmatrix} | & (-1)^2 \end{pmatrix}$

 V_0 , the correct plot of the probability density $(|\psi(x)|^2)$ with distance (x) is



- 107. A mixture of CaO and CaCO₃ is analyzed using thermogravimetry (TG) technique. The TG curve of the sample indicates that there is a mass change from 155.2 mg to 125.3 mg. The percentage of CaCO₃ in the mixture is close to
 (a) 54.2%(b) 27.1%(c) 43.8%(d) 29.9%
- 108. Match the given metal species in List I with the corresponding properties in List II List-II List-II

(A) $[Co(H_2O)_6]^{2+}$
(B) $[Cr(H_2O)_6^{-13+}$
(C) $NiCl_2(PPh_3)_2$
(D) $Pd(PPh_3)_4$ (I) magnetic moment higher than spin-only value and weak JT distortion
(II) spin-only magnetic moment and absence of JT distortion
(III) paramagnetic and tetrahedral
(IV) diamagnetic and tetrahedral
(IV

and $\langle p_x^2 \rangle = m\hbar\omega(n+1/2)$ for the quantum number *n*. The product of uncertainty of position and momentum for n = 1 is (a) $3\hbar/2$ (b) $\hbar/2$ (c) $2\hbar$ (d) \hbar

110. The variation of molar heat capacity at constant volume $(C_{v,m})$ with temperature (T) of a gaseous diatomic molecule is shown in the diagram below. The values of **X**, **Y** and **Z**, respectively, are [the diagram is not to the scale and discontinuity in the diagram represents dissociation]





111.	At 25°C, the total volume	e, V (in cm ³) of an ethan	nol solution containing	ng 1.0 kg of water fits the
	following expression V/c	$cm^3 = 1000 + 60\left(\frac{m}{m^0}\right) - 0$	$0.5 \left(\frac{m}{m^0}\right)^2$. Here <i>m</i> is	s molality and $m^0 = 1$ mol
	kg ⁻¹ . The partial molar ve	olume of ethanol (in cm ³	mol ⁻¹) in the solution	on prepared by mixing 460
	g of ethanol and 2 kg of	water is [Molar mass c	of ethanol = 46 g m	b^{1}]
	(a) 50	(b) 40	(c) 55	(d) 45

112. List I and List II give the molecular formula and the geometry of the species, respectively. List-I List-II

- (A) $\left[Zn \left\{ N \left(CH_2 CH_2 NH_2 \right)_3 \right\} Cl \right]^+$ (I) Trigonal bipyramidal (B) $\left[Cu \left(2, 2 - bpy \right) \left\{ NH \left(CH_2 COO \right)_2 \right\} \right]$ (II) Square pyramidal
- (C) $[ZrF_7]^{3-}$
- (D) $\left[AgTe_7 \right]^{3-}$ (IV) Trigonal planar

The option showing the correct match of species in List I and metal geometry in List II is(a) A-I, B-II, C-III, D-IV(b) A-I, B-II, C-IV, D-III(c) A-I, B-III, C-II, D-IV(d) A-II, B-I, C-III, D-IV

(III) Monocapped trigonal prism

- 113. Assuming the molecules are static, the ¹⁹F NMR spectra of $CIF_3(X)$ and $CIF_5(Y)$ consists of
 - (a) X: doublet and triplet; Y: singlet
 - (b) X: singlet; Y: singlet
 - (c) X: doublet and triplet; Y: doublet and quintet
 - (d) X: singlet; Y: triplet and quartet
- 114. The correct match of the labelled protons for ethyl vinyl ether in **Column P** with their chemical shift in **Column Q** is

	I ID		
	H _a H _c		
	ÓEt		
Column P	Colur	nn Q	
(A) H_a	(I) 6.4	45 (dd, $J = 13$, 7 Hz)	
(B) H_{b}	(II) 4	.05 (dd, J = 7, 2 Hz)	
(C) H_c	(III) 4	4.20 (dd, $J = 13, 2$ Hz	
(a) A-I, B-III, C-II	(b) A-I, B-II, C-III	(c) A-III, B-I, C-II	(d) A-II, B-I, C-III
The correct sequenc	e of reagents to effec	t the following transfo	ormation is
<u>∽s</u>	\blacktriangleright	СНО	

- (a) i. heat; ii. n-BuLi; iii. n-BuBr; iv. HgCl₂, H₂O
- (b) i. n-BuLi; ii. n-BuBr; iii. heat; iv. HgCl,, H₂O
- (c) i. heat; ii. HgCl₂, H₂O; iii. n-BuLi; iv. n-BuBr
- (d) i. n-BuLi; ii. n-BuBr; iii. HgCl₂, H₂O; iv. Heat

115.



- 116. To an aqueous solution of NaX and NaY, the addition of sulphamic acid (H_2NSO_3H) followed by acidification releases a nitrogen-containing gas P. Addition of Kl and a starch solution does not yield a blue color, indicating complete removal of NaX. However, the blue color appears when a piece of granulated Zn is added. The reaction proceeds with the evolution of a nitrogen containing gas Q. The correct option of **X**, **Y**, **P** and **Q**, respectively, is (a) $[NO_2]^-, [NO_3]^-, N_2, NO$ (b) $[NO_3]^-, [NO_2]^-, N_2, NO_3$ (c) $[NO_3]^-, [NO_2]^-, NO, N_2$ (d) $[NO_2]^-, [NO_3]^-, NO, N_2$ 117. Consider the following statements regarding the electronic spectra of lanthanide complexes. **A.** They exhibit fewer absorption bands due to a small number of microstates **B.** Their spectra are dependent on coordination number and geometry C. Molar extinction coefficients (e) are smaller compared to transition metal complexes **D.** Their absorption bands are sharp due to weak vibronic coupling E. Ligand field effects are negligible The option with correct statements is (a) A, B, D only (b) A, C, D only (c) C, D, E only (d) B, C, E only 118. Consider the following reactions (A) $SbCl_5 + AlCl_3 \longrightarrow [AlCl_4]^{-} [SbCl_4]^{+}$ (B) $AsF_3 + SbF_5 \longrightarrow [AsF_2]^+ [SbF_6]^-$ (C) NOF+SbF₅ \longrightarrow [NO]⁺[SbF₆]⁻ (D) $HF + SbF_5 \longrightarrow [SbF_4]^+ [HF_2]^-$ The correct option is (c) C and D only (d) A and D only (b) B and C only (a) A and B only 119. Cu crystallizes in face-centred cubic lattice. Considering each Cu atom as hard sphere and in contact with its nearest neighbors, the fraction of volume of the unit cell occupied by Cu atoms is (a) 0.16 (b) 0.56 (c) 0.74(d) 0.36 A molybdenum compound A is obtained by the CO displacement reaction of Mo(CO)₆ with PⁱPr₃, 120. reacts with H₂ to give compound B. The compounds A and B are (a) $\left[Mo(P^{i}Pr_{3})_{6} \right]$ and $\left[Mo(P^{i}Pr_{3})_{5}(\eta^{2}-H_{2}) \right]$ (b) $\left[Mo(CO)_3(P^iPr_3)_2 \right]$ and $\left[Mo(CO)_3(P^iPr_3)_2(\eta^2-H_2) \right]$ (c) $\left[Mo(CO)_3 (P^i Pr_3)_3 \right]$ and $\left[Mo(CO)_3 (P^i Pr_3)_2 (\eta^2 - H_2) \right]$
 - $(d) \left[Mo(CO)_4 \left(P^i Pr_3 \right)_2 \right] \text{ and } \left[Mo(CO)_4 \left(P^i Pr_3 \right)_2 \left(H \right)_2 \right]$